



Age	Appointment Expectations				
Newborn Check	Naked Weight / Length Head Circumference	Hep B @ birth			
1 Month Check	Naked Weight / Length Head Circumference				
2 Month Check	Naked Weight / Length Head Circumference	PENTACEL (DTap, HIB, IPV)	Hep B	PCV	ROTA* (oral)
4 Month Check	Naked Weight / Length Head Circumference	PENTACEL (DTap, HIB, IPV)		PCV	ROTA* (oral)
6 Month Check	Naked Weight / Length Head Circumference	PENTACEL (DTap, HIB, IPV)	Hep B	PCV	ROTA* (oral)
9 Month Check	Naked Weight / Length Head Circumference HCT/Lead Screening				
12 Month Check	Naked Weight / Length Head Circumference / TB screening/HCT & Lead level	MMR	Varicella	PCV	HepA
15 Month Check	Naked Weight / Length Head Circumference	DTap		HIB	
18 Month Check	Naked Weight / Length Head Circumference / Autism, TB & Lead Screening				HepA
2 Year	Weight / Height/BMI Head Circumference / HCT Autism, TB & Lead level				
3 Year	Weight / Height BP and Pulse / BMI TB & Lead Screening / HCT				
4 Year	Weight / Height / BMI - B/P Pulse Hearing & Vision Screening TB & Lead Screen/, HCT	MMR	Varicella	.PCV13^ (Booster)	Kinrix (DTap, IPV)
5-10 Years	Weight / Height / BMI - B/P Pulse Hearing & Vision Screening / TB & Lead Risk Screening HCT				
11-13 Years	Weight / Height / BMI - B/P Pulse Hearing & Vision Screening / TB & HCT	Boostrix (Tdap)	Menactra (Mcv4)		Gardasil X 3 @ 0, 2, & 6 mos
14-17 Years	Weight / Height / BMI - B/P Pulse Hearing & Vision Screening / TB & HCT				

---

<b>18</b> And Over	Weight / Height / BMI - B/P Pulse Hearing & Vision Screening / TB & HCT	
-----------------------	---	--

---

\*Rota Teq 3 doses and Rotarix 2 doses

^ PCV 13 booster for pt under 5yrs that haven't received the 13 serotypes

- Anyone 10 yrs or older must remain in office 15 minutes after any injection
- Patients receiving antibiotic injection must remain in office 20 minutes post injection